

# A Structural Interpretation of the 3D Poincaré Conjecture via the $n=2$ Uniqueness Theorem and Perelman's Ricci Flow

— Uniformization of Simply Connected Closed 3-Manifolds toward the Sphere, and a Structural Correspondence with Perelman's W-Entropy Functional — [Paper 07]

Yuuki YAMAGISHI (山岸 勇輝)

Yamagishi Complexity Torus Lab, Japan

September 2026 (3rd Edition)

— *From thinking to syncing.* —

## Abstract

For the three-dimensional Poincaré conjecture — “every simply connected closed 3-manifold is homeomorphic to the 3-sphere  $S^3$ ” (Perelman's theorem) — we present a structural interpretation based on the  $n = 2$  uniqueness theorem ( $dm/dr = 4\pi$ ). The argument proceeds in three stages.

**Stage one:** we set side by side the three conditions involved in the proof — the uniformity of shell mass  $n = 2$ , the Ricci flow equation  $\partial_t g_{ij} = -2R_{ij}$ , and simple connectivity  $\pi_1(M^3) = 0$  — and fix the logical position of each. What drives the convergence is the Ricci flow together with simple connectivity; the uniformity  $n = 2$  expresses, in the vocabulary of geometry, **a property satisfied by the destination  $S^3$**  (Section 2).

**Stage two:** we read the monotone increase of Perelman's  $\mathcal{W}$ -entropy functional  $\mathcal{W}(g, f, \tau)$  as the optimisation of rendering load in the Universe OS (Proposition 3.1). The  $4\pi$  in the factor  $(4\pi\tau)^{-3/2}$  of its integrand comes from the normalisation of the heat kernel, and its derivation runs along a different path from the solid angle  $\text{Vol}(S^2) = 4\pi$  — **the same number, at a different address** (Section 3).

**Stage three:** we place the uniformisation by the Ricci flow  $\partial_t g_{ij} = -2R_{ij}$  and surgery within the row formed by the resolution of the Poincaré conjecture in each dimension and the shell-conservation exponent  $n = d - 1$ , confirm that there is no derivation between the two, and give a dimension-by-dimension map (Section 4).

**Scope of this paper (scope note):** This paper presents a **structural interpretation** of a proven theorem. The three-dimensional Poincaré conjecture was proved by Perelman (2002–2003) using Hamilton’s Ricci flow and surgery; this paper takes no part in that proof and proposes neither an alternative proof nor a simplification. The monotonicity of the  $\mathcal{W}$  functional is likewise Perelman’s theorem and holds independently of the reading given here. Reading the Ricci flow as the optimisation of rendering load is an interpretation, not a theorem. The three conditions of Section 2 are not equivalent to one another; they are statements about different objects (a density distribution, a metric, a fundamental group). The Ricci flow used here is also, by Friedan’s theorem, the renormalisation group flow of the target metric of a two-dimensional sigma model (Paper 110). **What this paper adds** is that it expresses the uniformity of the destination  $\mathcal{S}^3$  of the Poincaré conjecture in the vocabulary of the  $n = 2$  uniqueness theorem; that it builds a table of correspondences between Perelman’s geometric concepts and the Universe OS picture; and that it tells apart the two different addresses — the normalisation of the heat kernel and the solid angle — from which the same  $4\pi$  appears.

## 1. Introduction: The Poincaré Conjecture and Perelman’s Geometrization Theorem

---

### 1.1 History of the 3D Poincaré Conjecture

The Poincaré conjecture, posed by Henri Poincaré (1904), was proved by Grigori Perelman (2002–2003) using Hamilton’s theory of Ricci flow together with surgery on singularities.

### 1.2 Positioning of the Structural Interpretation

This paper applies the  $n = 2$  uniqueness theorem (Paper 01) and the  $4\pi$  mass cancellation theorem (Paper 02) to Perelman’s mathematical proof, and offers the information-theoretic view that “the Ricci flow can be read as the spatial-memory optimisation and rendering-uniformisation process of the Universe OS.”

## 2. The Three Conditions (Triple Enforcement Principle) and Their Logical Position

---

### Proposition 2.1 (The Three Conditions and Their Logical Status)

Three conditions are involved in the process by which a simply connected closed 3-manifold  $M^3$  converges to the standard sphere  $S^3$ :

- i. **Analytic condition:** the uniformity condition for the shell mass growth rate of a spherically symmetric density distribution,  $n = 2$  ( $m(R) = 4\pi R$ ).
- ii. **Dynamical condition:** Perelman's Ricci flow equation  $\frac{\partial g_{ij}}{\partial t} = -2R_{ij}$ .
- iii. **Topological condition:** simple connectivity  $\pi_1(M^3) = 0$ .

The three conditions are statements about different objects. (i) is an analytic condition on a density distribution on  $\mathbb{R}^3$ , (ii) is a partial differential equation on a metric, and (iii) is a topological condition on the fundamental group; they are not equivalent to one another.

**The logical relation is as follows.** What actually works in the proof of the Poincaré conjecture is (ii) and (iii), due to Perelman (2002–2003). Condition (i) is **a property satisfied by the limit  $S^3$** , not a cause that drives the convergence. The role of (i) in this paper is **to supply a geometric vocabulary for characterising the destination**.

### 2.1 Structural Correspondence with Perelman's Proof

The correspondence between Perelman's geometric concepts and the information-geometric concepts of this framework is as follows:

- **Ricci flow**  $\partial_t g_{ij} = -2R_{ij} \iff$  smoothing protocol for spatial rendering load
- **Perelman's  $\mathcal{W}$ -entropy**  $\iff$  spatial memory compression efficiency (crash-avoidance rate)
- **Neck-pinch singularity**  $\iff n > 2$  micro-singularity freeze (data congestion)
- **Surgery**  $\iff$  forced reformatting by the Yamagishi patch  $m = 4\pi$
- **Standard sphere  $S^3$**   $\iff$  perfectly uniform rendering grid ( $m(R) = 4\pi R$ )

## 3. Structural Correspondence with Perelman's $\mathcal{W}$ -Entropy Functional

---

Perelman's  $\mathcal{W}$ -entropy functional is defined by

$$\mathcal{W}(g, f, \tau) = \int_{M^3} [\tau(R + |\nabla f|^2) + f - 3] (4\pi\tau)^{-3/2} e^{-f} dV$$

and increases monotonically under the Ricci flow ( $\frac{d\mathcal{W}}{dt} \geq 0$ ).

### Proposition 3.1 (Information-Theoretic Reading of $\mathcal{W}$ -Monotonicity)

The monotone increase  $\frac{d\mathcal{W}}{dt} \geq 0$  of the  $\mathcal{W}$  functional expresses the process by which the Ricci flow smooths out irregularities of curvature and drives the manifold toward a uniform state. Read in the Universe OS picture, it can be associated with **an optimisation protocol minimising the spatial rendering load**. Perelman himself introduced  $\mathcal{W}$  by analogy with entropy in statistical mechanics.

### 3.1 Two Addresses of the Factor $4\pi$

The  $4\pi$  in the factor  $(4\pi\tau)^{-3/2}$  of the integrand and the solid angle of the two-sphere  $\text{Vol}(S^2) = 4\pi$  are numerically equal, but their derivations run along different paths.

- The  $4\pi$  in  $(4\pi\tau)^{-3/2}$  is **the normalisation constant of the heat kernel (Gaussian kernel)**. It arises from the one-dimensional Gaussian integral

$$\int_{-\infty}^{\infty} e^{-x^2/(4\tau)} dx = \sqrt{4\pi\tau}$$

and in three dimensions is cubed to give  $(4\pi\tau)^{3/2}$ . The  $4$  here comes from the convention  $e^{-|x|^2/(4\tau)}$  in the exponent for the heat equation  $\partial_t u = \Delta u$ , and the  $\pi$  from the Gaussian integral  $\int e^{-x^2} dx = \sqrt{\pi}$ .

- $\text{Vol}(S^2) = 4\pi$  is **the solid angle of the two-sphere**, arising from  $\int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\phi = 4\pi$ .

**The two agree numerically, but their derivations share nothing.** The discipline of Paper 48 — not to count numerical coincidences of different origin as one — applies here too.

The monotonicity of the  $\mathcal{W}$  functional itself is a rigorous theorem of Perelman (2002) and holds independently of the interpretation in this paper.

## 4. Perelman's Theorem and an Arrangement by Dimension

**Theorem 4.1 (Perelman 2002–2003):** A simply connected closed 3-manifold  $M^3$  is homeomorphic to the standard 3-sphere  $S^3$ .

Theorem 4.1 is Perelman's theorem, and its proof rests on the Ricci flow and surgery. This paper gives a reading of that theorem.

4.1 Arrangement by Dimension

We set side by side the resolution of the Poincaré conjecture in each dimension and the corresponding shell-conservation exponent. The relation between the exponent  $n$  and  $\mathbf{Vol}(S^{d-1})$  is Theorem 4.1 of Paper 06 ( $n = d - 1$ ,  $\mathbf{Vol}(S^{d-1}) = 2\pi^{d/2}/\Gamma(d/2)$ ).

| Dimension $d$ | Resolution                   | $n = d - 1$ | $\mathbf{Vol}(S^{d-1})$  |
|---------------|------------------------------|-------------|--------------------------|
| 2             | Classical                    | $n = 1$     | $2\pi$                   |
| 3             | Perelman (2002–2003)         | $n = 2$     | $4\pi$                   |
| 4             | Freedman (1982), topological | $n = 3$     | $2\pi^2$                 |
| $\geq 5$      | Smale (1961)                 | $n = d - 1$ | $2\pi^{d/2}/\Gamma(d/2)$ |

There is no derivation between the right two columns and the left two; the table is **for setting the two side by side**. That the higher dimensions were settled earlier (Smale 1961 → Freedman 1982 → Perelman 2003) reflects the topological fact that surgery has more room in higher dimensions, and is a different matter from the values of  $\mathbf{Vol}(S^{d-1})$ .

5. Conclusion

This paper has presented a structural interpretation of the three-dimensional Poincaré conjecture proved by Perelman.

1. The logical position of the three conditions involved in the proof: what drives the convergence is the Ricci flow together with simple connectivity, and the uniformity  $n = 2$  expresses a property of the destination  $S^3$  (Section 2).
2. A view that reads the monotone increase of the  $\mathcal{W}$  functional as the optimisation of rendering load (Proposition 3.1).
3. The  $4\pi$  of the heat kernel and the  $4\pi$  of the solid angle are the same number at two different addresses (Section 3).
4. The resolution of the Poincaré conjecture in each dimension set side by side with the shell-conservation exponent (Section 4).

**Closing remark:** the view that reads uniformisation by the Ricci flow as “optimisation of rendering load” gives an intuitive entrance to this deep mathematical result. And just inside that entrance, the same  $4\pi$  appears from two different places: the spreading of heat and the angle over all directions.

# Revision Record

---

**3rd edition (September 2026):** The format was brought into line with the current template (the positioning and the relation to Paper 110 gathered into the scope note directly after the abstract, and the closing “statement of scope” removed). In the title, “Information-Geometric Reformulation” became “A Structural Interpretation”; in the subtitle, “Spherical Enforcement” and “Isomorphism” became “Uniformization toward the Sphere” and “Structural Correspondence”; and the abstract’s “we prove that it is isomorphic” was brought into line with the body as the presentation of a reading. DOIs were added to the corpus references.

**2nd edition (July 2026):** Three statements of the first edition were removed — that the three conditions are equivalent to one another, that the  $4\pi$  in  $(4\pi r)^{-3/2}$  is the solid angle  $\text{Vol}(S^2)$ , and the presentation as though this paper proved the main theorem.

# References

---

1. Poincaré, H. (1904). *Cinquième complément à l'analysis situs*. Rendiconti del Circolo Matematico di Palermo.

2. Perelman, G. (2002). *The entropy formula for the Ricci flow and its geometric applications*. arXiv:math/0211159.

3. Perelman, G. (2003). *Ricci flow with surgery on three-manifolds*. arXiv:math/0303109.

4. Hamilton, R. S. (1982). *Three-manifolds with positive Ricci curvature*. Journal of Differential Geometry.

5. Yamagishi, Y. (2026). *The Uniqueness Theorem of  $\mathfrak{n} = 2$  for Spherically Symmetric Density Distributions* [Paper 01]. Zenodo. 10.5281/zenodo.21403186

6. Yamagishi, Y. (2026). *The  $4\pi$  Mass Cancellation Theorem* [Paper 02]. Zenodo. 10.5281/zenodo.21403931

7. Yamagishi, Y. (2026). *The Critical Line and the Fixed Point of the Functional Equation* [Paper 06]. Zenodo. 10.5281/zenodo.21404262

8. Yamagishi, Y. (2026). *The Pfaffian's  $2^{\mathfrak{n}}\mathfrak{n}!$  Is One Block* [Paper 48]. Zenodo. 10.5281/zenodo.21602525

9. Yamagishi, Y. (2026). *Ricci Flow Is a Renormalization Group Flow* [Paper 110]. Zenodo. 10.5281/zenodo.21837476

— From thinking to syncing. —

Yuuki YAMAGISHI (山岸 勇輝)